#### Eastern Indian Ocean Upwelling Research Initiative (EIOURI)


As IIOE-2/YMC overlapping interest

Weidong Yu, First Institute of Oceanography, SOA, China




With contributions from JAMSTEC, UT, WHOI, Scripps, UM, AMFR, LIPI, BMKG ...


## Seasonal Upwelling Regions in Indian Ocean



SST (color) Wind Velocity in the Southwest Monsoon



# Research Moored Array for African-Asian-Australian Monsoon Analysis and Prediction (RAMA)



## Eastern Indian Ocean Upwelling Research Initiative (EIOURI) Science Plan and Implementation Strategy Writing Group Meeting


9-12 Apr. 2014, Phuket, Thailand



Hosted by: Thailand-China Joint Laboratory for Climate and Marine Ecosystem

- Physical oceanography
  - Upwelling processes: local and remote forcing
  - Open ocean-coastal interactions: circulation and eddy
- BGC and Biological oceanography
  - Nutrient and plankton responses (e.g. classic vs cryptic upwelling, stoichiometry)
- Ecology
  - Food web, higher level responses and physical mechanisms (e.g. offshore advection, massive-kills)
- The human dimension

## **Pilot Cruise in 2013**





# Extension of EIOURI to YMC in collaboration with MOMSEI

- Indonesian-Australian Monsoon Onset
- In context of MOMSEI study in BoB
- To identify the first eastward propagating MJO which triggers Indonesian-Australian Monsoon


### **Monsoon Onset Process over BoB**


(SST-MJO-Monsoon Onset)





### **Composite of BoB Monsoon Onset**








Fig. 10. Composite plan view of OLR relative to onset at Darwin. Contour interval is 5 W  $m^{-2}$  with first contour at  $\pm 10$  W  $m^{-2}$ . The sequence is (a) day -8, (b) day +2, and (c) day +12.

Monsoon onset coincides with the first occurrence of convectively active 40-50 day oscillation.

Again, SST-MJO-Onset relationship should be examined, as in BoB.

